

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
17 January 2002 (17.01.2002)

PCT

(10) International Publication Number  
**WO 02/04681 A2**

(51) International Patent Classification<sup>7</sup>: C12Q 1/68

(21) International Application Number: PCT/US01/21846

(22) International Filing Date: 11 July 2001 (11.07.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/217,782 11 July 2000 (11.07.2000) US

(71) Applicant: NANOSPHERRE INC. [US/US]; 1801  
Maple Avenue, Evanston, IL (US).

(72) Inventors: LETSINGER, Robert, L.; 316 Third Street,  
Wilmette, IL 60091 (US). GARIMELLA, Viswanadham;  
1915 Maple Avenue, Englehart Hall, Apt. 323, Evanston,  
IL 60201 (US).

(74) Agent: MIAO, Emily; McDonnell Boehnen Hulbert &  
Berghoff, 300 South Wacker Drive, Suite 3200, Chicago,  
IL 60606 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,  
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,  
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,  
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,  
TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished  
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

WO 02/04681 A2

(54) Title: METHOD OF DETECTION BY ENHANCEMENT OF SILVER STAINING

(57) Abstract: The present invention relates to a method for amplifying a detection signal by enhancing or promoting the deposition of additional silver in assay detection systems where the formation of a silver spot serves as a reporter for the presence of a target molecule, including biological polymers (e.g., proteins and nucleic acids) and small molecules.

**THIS PAGE BLANK (USPTO)**

**METHOD OF DETECTION BY ENHANCEMENT OF SILVER STAINING****CROSS-REFERENCE:**

- 5 This application claims the benefit of priority from U.S. Provisional application No. 60/217,782, filed July 11, 2000.

**FIELD OF THE INVENTION:**

- The invention relates to several fields, especially to the detection of specific  
10 nucleic acids, proteins, carbohydrates, or organic compounds immobilized on a solid surface. More particularly it relates to detection systems in which the immobilized target is recognized by a metallic nanoparticle probe and for which the signal for detection has been amplified by reductive deposition of silver on the nanoparticle probe.

15 **BACKGROUND OF THE INVENTION:**

(a) *Gold Nanoparticle Probes:*

- The use of gold nanoparticle probes as reporter for detection of biological polymers was first described by W.P. Faulk and G.M. Taylor, who employed the nanoparticles as immunocytochemical probes for surface antigens [*Immunochemistry*, 8,  
20 1081 (1971)]. Since then gold colloids have been widely used for detection of a variety of proteins using electron or light-microscopy to observe the particles [for reviews see Hacker, G. W. in *Colloidal Gold; Principles, Methods, and Applications*, Vol. 1, Academic Press, Inc. (1998) p 297, and Garzon, S., and Bendayan, M. in *Immuno-Gold Electron Microscopy in Virus Diagnosis and Research*, Ed. Hyatt, A.D. and Eaton, B.T.,  
25 CRC Press, Ann Arbor, (1993) p 137]. Recently, applications of gold nanoparticle or cluster conjugates as probes for detection of oligonucleotides and nucleic acids have been suggested [Kidwell, D.A., and Conyers, S.M., United States Patent 5,384,265 (1995);

-2-

Hainfeld, J.F., et al. United States Patent 5,521,289 (1996)] and described [Tomlinson, S., et al., *Analytical Biochemistry*, 171, 217 (1988); Mirkin et al., *Nature*, 15, 607 (1996); Storhoff, J.J. et al., *J. Am. Chem. Soc.*, 120, 1959 (1998)].

**(b) Silver Enhancement of Signal:**

5 It has been found that the sensitivity for assays utilizing gold markers for proteins in tissues [Danscher, G. *Histochemistry*, 71, 1 (1981); Holgate, C.S. et al. *J. Histochem. Cytochem.* 31, 938 (1983)], for nucleic acids visualized *in situ* in immobilized biological systems [Gassell, G.J., et al., *J. Cell Biology*, 126, 863 (1994); Zehbe, I. et al., *Am J. of Pathology*, 150, 1553 (1997); Hacker, G. W., *Eur. J. Histochem* 42, 111 (1993) and for  
10 gold probes targeted to oligonucleotides captured on oligonucleotide arrays on a glass surface [T. A. Taton, C.A. Mirkin, R.L. Letsinger, *Science*, 289, 1757 (2000)] can be significantly increased by silver staining. In this process, the gold particles captured on a surface are treated with a solution containing silver ions and a reducing agent (e.g., hydroquinone). The gold catalyzes reduction of the silver ions so that silver is deposited  
15 on the gold particle, and the early-deposited silver can itself catalyze further reduction of silver ion. As a consequence, the amount of metal that can be visualized is greatly increased. Unfortunately, however, the silver reduction catalyzed by the deposited silver ceases after a time, so the extent of amplification achievable is limited. When employed in enhancing visibility of gold nanoparticles on a glass plate, one observes darkening of  
20 the spot characteristic for the gold probes captured by a target sequence. Indeed, a good silver spot may be observed for cases where the amount of gold deposited initially is too small to be visible to the naked eye. Typically, the reaction time for the silver staining step is short, of the order of five minutes or less. Long exposure to the silver solution leads to non-selective deposition of silver metal and a high background. The silver ion

-3-

solution and the reducing agent are mixed just prior to application to minimize the uncatalyzed reduction.

(c) *Oligo- and Polynucleotide Arrays:*

A recent major innovation in biology utilizes arrays of oligonucleotides or polynucleotides tethered to a solid surface. These oligomers serve as capture probes to bind complementary DNA or RNA target sequences. The captured sequences can in turn be recognized by fluorescent labels previously attached to them or by fluorescent or colorimetric probes that bind to a segment of the target. As stated by Eric Lander [*Nature Genetics Supplement*, 21, 3 (1999)]: "Arrays offer the first great hope....by providing a systematic way to survey DNA and RNA variation. They seem likely to become a standard tool of both molecular biology research and clinical diagnostics. These prospects have attracted great interest and investment from both the public and private sectors."

Array technology is indeed now greatly accelerating developments in our understanding of genetic variation and gene expression. Nevertheless, current methodology suffers from several limitations, an important one being relatively low sensitivity in detecting fluorescently labeled targets on the chip arrays. Typically, targets in the range of picomolar concentrations or higher must be employed. Genetic analyses of natural targets in the attomolar or zeptomolar range therefore require target amplification by PCR. This procedure demands time and labor, and the target amplification can lead to errors in the sequence to be tested.

A need exists for a more sensitive, simpler, and cheaper detection method for polynucleotides arrayed on chips. Progress in detection technology has been made with the use of gold nanoparticle oligonucleotide conjugates as probes and signal amplification by silver ion reduction, which enables assays of polynucleotides of 50 fM concentration to be readily detected [for the methodology, see T. A. Taton, C.A. Mirkin, R.L. Letsinger,

-4-

*Science*, 289, 1757 (2000). We describe here a discovery that significantly lowers further the target concentration required for assays employing gold nanoparticles and other metallic nanoparticles.

## 5 SUMMARY OF THE INVENTION:

The present invention relates to a method for amplifying signal by enhancing the deposition of silver in detecting systems where the formation of a silver spot serves as a reporter for the presence of a molecule, including biological polymers (*e.g.*, proteins and nucleic acids) and small molecules. The detecting systems include detection of molecules  
10 *in situ* (*e.g.*, on cells or in a tissue sample) and assays where the molecule to be detected (the target molecule) is bound to a substrate or is captured by another molecule bound to a substrate (the capture molecule). The invention has special utility in increasing the signal strength in diagnostic and screening applications involving detection of target molecules arrayed at discrete positions on a solid surface. It, therefore, provides a means for greatly  
15 enhancing the sensitivity of tests carried out on microarrays or microchips. The process is distinguished by the simplicity and economy of its execution and the large enhancement in signal and, thereby, sensitivity realized.

This invention is based on the discoveries that (1) gold nanoparticles coated with oligonucleotides bind to silver that had previously been deposited on gold nanoparticle-  
20 oligonucleotide conjugates immobilized by hybridization on a glass substrate or plate and (2) that the resulting (gold nanoparticle-oligonucleotide-silver-(gold-oligonucleotide) structures function as catalyst for the further deposition of silver by reduction of silver ions. The first discovery is surprising since one might expect that the surface bound oligonucleotides, which shield the nanoparticles from non-specific binding to the glass  
25 surface and the oligonucleotides immobilized on the glass surface, would also shield the

-5-

nanoparticles against interaction with the silver surface. Indeed, other work has shown that oligonucleotides protect gold nanoparticle oligonucleotide conjugates from fusing to form gold-gold bonds between individual nanoparticles even when the mixtures are dried. The second discovery is significant since it provides a means for substantially increasing the metallic mass at the site of the originally immobilized nanoparticles. In conjunction with development of buffer conditions that enable oligonucleotide nanoparticle conjugates that are unbound by hybridization or interaction with silver to be washed cleanly from the glass surface these findings opened opportunities for assaying polynucleotide targets at extremely low target concentrations.

10 Accordingly, one objective of the invention is to provide a method for amplifying a detection signal comprising:

- (a) providing a substrate having deposited silver;
- (b) contacting the substrate having deposited silver with a solution comprising nanoparticles having oligonucleotides bound thereto so as to produce a substrate having a nanoparticles-silver sandwich;
- 15 (c) washing the substrate having said sandwich; and
- (d) contacting the substrate having said sandwich with silver ions and a reducing agent to promote silver deposition onto the nanoparticles of said sandwich.

20 The nanoparticles having oligonucleotides bound thereto comprise gold, silver, platinum or mixtures thereof. These nanoparticles may be in the form of gold nanoparticle-oligonucleotide conjugates or complexes.

Another object of the invention is to provide a method for promoting silver deposition onto a surface comprising silver, said method comprising the steps of:

- (a) providing a surface having silver bound thereto;
- 25 (b) contacting the surface with a solution comprising nanoparticles having oligonucleotides bound thereto so as to produce a surface having a nanoparticles-silver sandwich ;
- (c) washing the surface having said nanoparticles-silver sandwich;

-6-

- (d) contacting the surface having said nanoparticles-silver sandwich with a solution including silver ions under reducing conditions to promote silver deposition onto said nanoparticles of said nanoparticles-silver sandwich; and
- (e) washing the surface having deposited silver.

5 According to this method, the surface may include cells or tissue for in situ detection of target molecules. Preferrably the nanoparticles having oligonucleotides bound thereto comprise gold nanoparticles having oligonucleotides bound thereto in conjugate or complex form. In practicing this invention, gold nanoparticle oligonucleotide conjugates are preferred.

10 A further object of the invention is to provide a kit for detection signal amplification comprising:

- (a) container including nanoparticles having oligonucleotides bound thereto;
- (b) container including a silver salt; and
- (c) container including a reducing agent.

15 The kit may include instructions for use in amplifying silver stain detection signals. Preferrably the nanoparticles having oligonucleotides bound thereto comprise gold nanoparticles having oligonucleotides bound thereto in conjugate or complex form. In practicing this invention, gold nanoparticle oligonucleotide conjugates are preferred.

## 20 DESCRIPTION OF THE FIGURES

Figure 1 is a schematic representation of amplification of a detection signal for a gold nanoparticle immobilized on a solid surface. Step (a) is conventional silver staining. In step (c), gold nanoparticles bearing oligonucleotides are bound to the silver surface. Step (e) is conventional silver staining of the gold nanoparticles that were bound to the

25 previous silver surface.

Figure 2 illustrates two plates with or without application of the amplification method of the invention. Plate (a) was subjected to a second round of conventional silver staining in an attempt to further enhance the spot. Slight but very little signal



-7-

enhancement took place. Plate (b) was subjected to the signal amplification method of the invention.

Figure 3 illustrates two plates with or without application of the amplification method of the invention. Plate (a) indicates the signal obtained for a sample using conventional silver staining. Plate (b) shows the enhancement achievable by the application of the inventive signal amplification method.

Figure 4 illustrates two plates with or without application of the amplification method of the invention. Plate (a) shows the results of conventional silver staining for a test in which a 25 fM solution of the target oligonucleotide was used. Plate (b) shows the results for the same test in which the initial silver staining was followed with the signal amplification method of the invention.

Figure 5 illustrates a plate following application of the inventive signal amplification method of the invention. Spot obtained after two cycles of the nanoparticle-oligonucleotide (conjugate I)/silver staining methodology applied when gold nanoparticle probes and an oligonucleotide target (63-mer) were immobilized on a glass plate. The concentration of the target oligonucleotide in this case was only 1 fM.

## DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to a method for signal amplification for detecting target substances. In particular, the invention provides for amplification of signal and enhancement of assay sensitivity for detection of minute quantities of target molecules, e.g., nucleic acids, that are based on silver stain detection. In practicing this invention, the inventive method is used after the application of conventional silver staining where it leads to the formation of silver-gold(oligonucleotide)-silver assemblies, termed here silver-gold'-silver sandwich assemblies.

-8-

The prior steps leading up to and including conventional silver staining can, for example, involve: capture of a target oligonucleotide by an oligonucleotide capture probe bound to a glass surface, washing, addition of nanoparticle oligonucleotide conjugates complementary to an unhybridized segment of the bound target oligonucleotide, washing, addition of a silver staining solution (Ag<sup>+</sup> plus a reducing agent, available through Sigma), washing, drying, and viewing, either with the naked eye or with aid of pictures obtained using a simple flat-bed scanner. This procedure was described by Taton et al., *Science*, 289, 1757 (2000), who showed that although the gold spot obtained was too weak to observe directly for target concentrations of 100 nM in oligonucleotide, conventional silver deposition affords a strong dark spot. The limit of detection for this system using conventional silver staining is 50 fM concentrations, for which an extremely faint spot is observed. Re-exposure to the silver solution did not noticeably further enhance the intensity of the spot.

For amplification of the silver signal by the inventive silver-gold'-silver sandwich method, the glass plate containing silver stain from a prior application of conventional staining is further exposed to an aqueous amplification solution of nanoparticles having oligonucleotides bound thereto, preferably gold nanoparticle-oligonucleotide conjugates or a gold-nanoparticle oligonucleotide complexes. The oligonucleotide sequence does not need to be related to the target oligonucleotide sequence or the sequences on the initial probes. For gold nanoparticle-oligonucleotide conjugates, the amplification solution generally contains between about 1 nM and about 5 nM nanoparticle oligonucleotide conjugates. The amplification solution may include salts, such as buffer salts, and detergents such as phosphate, citrate, HEPES, and NES and is preferably has a pH of about 7. In practicing the invention, an aqueous amplification solution including aqueous 0.1M NaCl and 10 mM phosphate (pH 7.0) was found to be particularly useful. For gold

-9-

nanoparticle oligonucleotide complexes, the amplification solution is the same. Any suitable mode of exposing the silver stain to the amplification solution may be used.

Representative examples include spraying, dipping, and the like.

After about 5 to 30 minutes, preferably about 10 minutes, the unbound  
5 nanoparticles conjugates or complexes are removed by washing the plate with a suitable aqueous solution, preferably water. Optionally, the plate is dried by any suitable method such as air drying. The plate is then re-exposed to a silver staining solution for a suitable time period, generally for about 5 to about 10 minutes, preferably about 5 minutes. It will  
10 be understood that any suitable silver staining method may be used. Suitable, but non-limiting, examples of useful silver staining methods and compositions are described, for instance, in M.A. Hayat, Ed., "Immunogold-Silver Staining," CRC Press (1995).

Generally, silver staining solutions include silver ion in the form of a silver salt such as silver acetate (preferred), silver lactate, and silver nitrate. These solutions also include a reducing agent that is admixed into the solution just prior to use. Suitable, but non-  
15 limiting, examples of reducing agents include hydroquinone, n-propyl galate, p-phenylenediamine, and formaldehyde. If desired, other agents such as gum Arabic may be used to mediate the silver stain process. Any suitable mode of contacting the substrate with the silver stain solution may be used. Representative examples include spraying, dipping, and the like.

20 Following washing with water (preferred) or other suitable solution (e.g., aqueous solution containing 0.1M NaCl and 10 mM phosphate (pH 7.0) to remove unreacted silver stain solution, the re-exposed plate is observed visually or copied by a flat-bed scanner. This treatment leads to a great increase in the darkness of the spot. The process may be repeated as often as desired to further enhance the amount of deposited silver and the  
25 darkness of the spot. With this inventive amplification method, one can readily observe a

-10-

dark silver spot for an assay using 25 fM target concentration. With two cycles of the new nanoparticle-silver sandwich procedure, 1 fM target solutions can be recognized, and with three cycles, 0.1 fM solutions give positive though weak spots. These experiments were carried out using 1 microliter of the target solution in each case. For the assay with 0.1 fM solution this corresponds to ~60 target molecules in the applied sample.

Any substrate can be used which allows observation of a silver stain as the detectable change. Suitable substrates include transparent solid surfaces (e.g., glass, quartz, plastics and other polymers. The substrate can be any shape or thickness, but generally will be flat and thin. Preferred are transparent substrates such as glass (e.g., glass slides) or plastics (e.g., wells of microtiter plates).

The silver stain signal amplification method of the invention depends on the use of nanoparticles-oligonucleotide conjugates or complexes that satisfy certain characteristics. First, the nanoparticles do not stick to the surface of the chip being tested. Ordinary nanoparticles prepared by the citrate reduction method of Frens (Frens, G., *Nat. Phys. Sci.*, 241, 20-22 (1973) are not satisfactory since they bind indiscriminately to the oligonucleotide-derivatized glass plate used as the substrate for these assays. Subsequent silver enhancement then gives false positives as dark areas. Second, the nanoparticles bind to a deposited silver surface such that on subsequent washing, the attached nanoparticles remain bound to the silver area while nanoparticles suspended in solution are cleanly removed. Third, the nanoparticles function as agents to reduce silver ions under silver staining conditions. In practicing this invention, useful nanoparticles are nanoparticles coated with oligonucleotides linked through sulfur to the surface (nanoparticle oligonucleotide conjugates) such as the ones described in J. J. Storhoff et al., *J. Am. Chem. Soc.*, 120, 1958 (1999) (for a specific example, see conjugate I in Example 1 below) or with natural type oligonucleotides adsorbed to the surface (nanoparticle

-11-

oligonucleotide complexes) such as conjugate III described in Example 1.

Both types of nanoparticles work well in low or moderate salt solution (e.g. up to 0.1 M), but the conjugates containing the sulfur anchor are particularly preferred for tests conducted at high salt concentration, at which the complexes formed by simple adsorption

5 of oligonucleotides are unstable and aggregate. It will be understood by the ordinary skilled artisan that any nanoparticle preparation that meets the criteria listed above are useful as intermediary agents in forming the sandwich assemblies, and the methodology can be applied for the amplification of the silver signal for any target visualized by an initial silver deposition. While gold nanoparticles are particularly preferred, any  
10 nanoparticle that catalyze the reduction of silver can be used including silver and platinum nanoparticles.

The preparation of nanoparticles suitable for use in the practice of the invention, the attachment of oligonucleotides to them, the flatbed scanner technique, and various assays formats for the detection of nucleic acids using conventional silver staining are  
15 described in co-pending applications Serial Nos. 09/760,500, filed January 12, 2001; 09/603,830, filed June 26, 2000; 09/344,667, filed June 25, 1999; 09/240,755, filed January 29, 1999; 60/031,809, filed July 29, 1996; 60/176,409; and 60/200,161, filed April 26, 2000; and international application Nos. PCT/US97/12783, filed July 21, 1997; PCT/US00/17507, filed June 26, 2000; and PCT/US01/01190, filed January 12, 2001,  
20 entitled "Nanoparticles Having Oligonucleotides Attached Thereto And Uses Therefor," the entire contents of which are incorporated herein by reference.

## EXAMPLES

### Example 1: Preparation of Oligonucleotide Modified Gold Nanoparticles.

Oligonucleotides and 5'-mercaptoalkyl-oligonucleotides were prepared using  
25 phosphoramidite chemistry as described by Storhoff et al. [*J. Am. Chem. Soc.* 120, 1959-

-12-

1964 (1998)]. Gold nanoparticles (~13 nm diameter) were prepared as described by Grabar, K.C., et al. [*J. Analyt. Chem.*, 67, 735-743 (1995); Frens, G., *Nat. Phys. Sci.*, 241, 20-22 (1973)].

For preparation of the nanoparticle conjugate (I) used in the sandwich silver-gold'-  
5 silver amplification scheme, 5'-mercaptoalkyl-oligonucleotide II was prepared and joined to gold nanoparticles by the general linking procedure described by Storhoff et al., [*J. Am. Chem. Soc.* 120, 1959-1964 (1998)]. The nanoparticle-oligonucleotide complex (III) was prepared by mixing 1  $\mu$ L of an aqueous solution containing 0.26 A<sub>260</sub> Units of  
10 oligonucleotide IV with 100  $\mu$ L of citrate stabilized gold colloid (prepared as described by Grabar et al. [*J. Analyt. Chem.*, 67, 735-743 (1995)]) and allowing the solution to stand overnight.

- I. Conjugate formed from gold nanoparticles and II.
- II. 5'-HS(CH<sub>2</sub>)<sub>6</sub>OP(O)(O<sub>2</sub>)O-(A)<sub>20</sub>TGGGTAGCAGACCTC (SEQ ID NO.:1)
- III. Complex formed from gold nanoparticles and IV
- 15 IV. 5'-GCTCTAGAATGAACGGTGAAGGCGGCAGG (SEQ ID NO.:2)

### Example 2: Sandwich Amplification of Silver Signal

In this Example, four separate experiments were conducted using the nanoparticle oligonucleotide conjugate (I) or complex (III) prepared as described in Example 1, and glass slides containing silver spots from oligonucleotide assays carried out using the silver  
20 staining method described by T.A. Taton et al, *Science*, 289, 1757-60 (2000). The glass plate bearing the silver spots was exposed to a solution of gold nanoparticle conjugate I (Figures 2, 4, and 5) or gold nanoparticle complex III (Figure 3) for 10 minutes, washed with 1 M NaNO<sub>3</sub> in nanopure water, and exposed for five minutes at room temperature to a 1:1 mixture of freshly mixed sample of the two commercial Silver Enhancer solutions  
25 (Catalog Nos. 55020 and 55145, Sigma Corporation, St. Louis, MO) for 5 minutes,

-13-

following the Sigma protocol for the silver staining step, including final washes with nanopure water, sodium thiosulfate solution, and nanopure water. The plate was dried and observed, both visually and by copying for records using a Hewlett Packard Scanner [Model no. 5200C]. The direct visual observations of spots corresponded to the prints  
5 obtained using the scanner. The figures are enlarged.

Figure 2 illustrates the results for the first experiment using nanoparticle oligonucleotide conjugate I. In this experiment, two plates, each containing very faint silver spots from previous hybridization and silver staining of a dilute target solution, were used. Plate (a) was subjected to a second round of conventional silver staining in an  
10 attempt to further enhance the spot. Slight but very little signal enhancement took place. Plate (b) was subjected to conditions for the gold sandwich amplification. This entailed exposing the plate containing the weak silver spots to the solution of nanoparticle conjugate I for 10 minutes, followed by washing with water and conventional silver staining for 4 minutes. The darkness of the spots in plate (b) relative to those in plate (a)  
15 clearly demonstrate the power of the new metal sandwich or signal enhancement method.

Figure 3 illustrates the results for an experiment using nanoparticle oligonucleotide complex III. Plate (a) indicates the signal obtained for a sample using conventional silver staining. Plate (b) shows the enhancement achievable by the gold sandwich technology. Note that two sample spots are shown in each case. For plate (b), a plate containing a  
20 weak silver spot corresponding to that in plate (a) was exposed to a solution of nanoparticle oligonucleotide complex III, and following washing, was subjected to conventional silver staining. In this case, three parts of colloid III, prepared as described above, was mixed with seven parts of 10 mM phosphate buffer at pH 7. The plate was then exposed to this colloid solution for 10 minutes, washed with nanopure water, and  
25 exposed to the silver staining mixture for 4 minutes as before.

-14-

Figure 4 illustrates the results for an experiment using gold nanoparticle oligonucleotide I to enhance the weak silver signal resulting from an assay of a 63 nucleotide target oligonucleotide

5' GTA GGC GAA.CCC TGC CCA GGT CGA CAC ATA GG T GAG GTC TGC TAC

5 CCA CAG CCG GTT AGG TGC 3' (SEQ ID NO.: 3)

at 25 fM concentration. Note that our standard assay using just silver amplification gives a very weak spot (plate (a)). Plate (a) shows the results of conventional silver staining for a test in which a 25 fM solution of the target oligonucleotide was used. Plate (b) shows the results for the same test in which the initial silver staining was following by 10 minute exposure to the colloidal solution of nanoparticle conjugate I, followed by silver staining. In this case, the gold solution had been diluted six-fold with 0.1 M NaCl-10 mM phosphate buffer. Note the strong signal for a target, captured at 25 fM concentration, after the metal sandwich enhancement procedure.

Figure 5 illustrates the results for an experiment using nanoparticle oligonucleotide conjugate I to enhance the silver signal obtained from an assay carried out using a 63-mer target [see SEQ ID NO.: 3] at a concentration of 1 fM. In this case, our standard procedure for silver staining failed to show any spot. Double enhancement using the gold-silver treatment, however, showed a strong signal, as shown. This experiment was carried out by treating the plate that had been exposed to silver amplification by the standard procedure successively with: (1) gold nanoparticle conjugate I (1.5 nM in nanoparticles in 0.1 M NaCl and 10 mM phosphate buffer at pH 7.0) for 10 minutes; (2) wash with 1 M NaNO<sub>3</sub>; (3) treat with a mixture of Silver Enhancer solutions (catalog nos. 5020 and S 5145, Sigma Corporation, by the Sigma protocol; (4) wash with 1 M sodium nitrate; (5) repeat steps 1-4; (6) wash with sodium thiosulfate ; and (7) wash with water.



-15-

**WHAT WE CLAIM:**

1. A method for amplifying a signal comprising:
  - (a) providing a substrate having deposited silver;
  - 5 (b) contacting the substrate having deposited silver with a solution comprising nanoparticles having oligonucleotides bound thereto so as to produce a substrate having a nanoparticles-silver sandwich;
  - (c) washing the substrate having said sandwich; and
  - (d) contacting the substrate having said sandwich with silver ions and a  
10 reducing agent to promote silver deposition onto the nanoparticles of said sandwich and produce silver-nanoparticles-silver sandwich.
  - (e) washing the substrate having the silver-nanoparticles-silver sandwich.
2. The method according to claim 1, wherein the nanoparticles comprise gold,  
15 silver, platinum or mixtures thereof.
3. The method according to claim 3, wherein the nanoparticles having oligonucleotides bound thereto comprise gold nanoparticle-oligonucleotide conjugates or complexes.  
20
4. The method according to claim 2, wherein the silver ion is derived from a silver salt comprising silver acetate, silver lactate, or silver nitrate.
5. The method according to claim 1 wherein the reducing agent comprises  
25 hydroquinone, n-propyl galate, p-phenylenediamine, or formaldehyde.
6. The method according to claim 1 wherein step (b) contacting is performed for a period ranging from about 5 to 30 minutes.
- 30 7. The method according to claim 6 wherein step (b) contacting is performed for about 10 minutes.
8. The method according to claim 1 wherein step (d) contacting is performed for a period ranging from about 5 to 10 minutes.

-16-

9. The method according to claim 8 wherein step (d) contacting is performed for about 5 minutes.

5 10. The method according to claim 1 wherein the deposited silver bound to the substrate is arranged in the form of an array.

11. The method according to claim 1, wherein step (c) washing is performed with water.

10 12. The method according to claim 1, wherein the substrate is glass.

13. A method for promoting silver deposition onto a surface comprising silver, said method comprising the steps of:

- 15 (a) providing a surface having silver bound thereto;
- (b) contacting the surface with a solution comprising nanoparticles having oligonucleotides bound thereto so as to produce a surface having a nanoparticles-silver sandwich ;
- (c) washing the surface having said nanoparticles-silver sandwich;
- 20 (d) contacting the surface having said nanoparticles-silver sandwich with a solution including silver ions under reducing conditions to promote silver deposition onto said nanoparticles of said nanoparticles-silver sandwich; and
- (e) washing the surface having deposited silver.

25 14. The method according to claim 13 wherein said surface comprises cells or tissue.

15. The method according to claim 13, wherein the nanoparticles comprise gold, silver, platinum or combinations thereof.

30

16. The method according to claim 13, wherein the nanoparticles having oligonucleotides bound thereto comprise gold nanoparticles having oligonucleotides bound thereto.

-17-

17. The method according to claim 16, wherein the gold nanoparticles having oligonucleotides bound thereto comprise gold nanoparticle- oligonucleotide conjugates or complexes.

5

18. The method according to claim 13, wherein the silver ion is derived from a silver salt comprising silver acetate, silver lactate, or silver nitrate.

19. The method according to claim 13 wherein the reducing agent comprising  
10 hydroquinone, n-propyl galate, p-phenylenediamine, or formaldehyde.

20. The method according to claim 13 wherein step (b) contacting is performed for a period ranging from about 5 to 30 minutes.

15 21. The method according to claim 20 wherein step (b) contacting is performed for a period of about 10 minutes.

22. The method according to claim 13 wherein step (d) contacting is performed for a period ranging from about 5 to 10 minutes.

20

23. The method according to claim 22 wherein step (d) contacting is performed for about 5 minutes.

24. The method according to claim 13 wherein the silver bound to the substrate  
25 are arranged in the form of an array.

25. A kit for signal amplification comprising:  
(b) container including nanoparticles having oligonucleotides bound thereto;  
(b) container including a silver salt; and  
30 (c) container including a reducing agent.

26. The kit according to claim 25, wherein the nanoparticles comprise gold, silver, platinum or combinations thereof.

-18-

27. The kit according to claim 26, wherein the nanoparticles comprising gold nanoparticles having oligonucleotides bound thereto.

5 28. The kit according to claim 27, wherein the gold nanoparticles having oligonucleotides bound thereto comprise gold nanoparticle-oligonucleotide conjugates or complexes.

29. The kit according to claim 25, wherein the silver ion is derived from a silver salt comprising silver acetate, silver lactate, or silver nitrate.

30. The kit according to claim 25, wherein the reducing agent comprises hydroquinone, n-propyl galate, p-phenylenediamine, or formaldehyde.




Figure 1. Schematic representation of amplification of a detection signal for a gold nanoparticle immobilized on a solid surface. Step "a" is conventional silver staining. In step "c" gold nanoparticles bearing oligonucleotides are bound to the silver surface. Step "e" is conventional silver staining of the gold nanoparticles that were bound to the previous silver surface.

**THIS PAGE BLANK (USPTO)**

Figure 2. (a) Silver signal after double silver staining.  
(b) Silver signal after one silver staining step followed by treatment with I and a second silver step.

(a)



(b)



**THIS PAGE BLANK (USPTO)**




Figure 3. (a) Two spots from a conventional silver staining step on a very dilute DNA sample

(b) Same spots after further development using nanoparticle-oligonucleotide-complex III.

(a)



(b)



**THIS PAGE BLANK (USPTO)**

Figure 4: Silver spots for assays that used 25fM 63-mer oligonucleotide target solutions.

- a) Spot from conventional silver staining.
- b) Spot for silver-gold-'silver procedure where the gold' reagent is conjugate I.

(a)




(b)



**THIS PAGE BLANK (USPTO)**

Figure 5: Silver spot obtained by amplifying the silver residue obtained by conventional silver staining for an assay of an oligonucleotide at 1 fM concentration, using nanoparticle probes. The initial silver spot was not visible. Amplification was achieved by two cycles of the gold nanoparticle(reagent I) / silver procedure.



**THIS PAGE BLANK (USPTO)**

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**